Antibiotic Medium B (Eur. Pharm.)

Art. No. 01-016

Also known as

Antibiotic Medium 10, Medium 10

Specification

Antibiotic Medium B is used in the microbiological assays of Colistimethate and Polymyxin by the Agar Diffusion method.

Formula* in g/L

Casein Pancreatic Digest.	17,00
Soybean meal Digest	3,00
Sodium chloride	5,00
Dextrose	2,50
Dipotassium phosphate	2,50
Agar	15,00
Final pH 7.3 ± 0.1 at 25°C	

Reconstitution

Add 45 g of powder to 1 L of distilled water. Once dissolved, add 10 mL of Polysorbate 80 (Art. No. TW0080). Bring to the boil and dispense into suitable containers. Sterilize in the autoclave at 121°C for 15 minutes.

Description

Antibiotic Medium B is recommended by the European Pharmacopoeia and the USP for determining antibiotic potency by the microbiological assay techniques, specifically for colistimethate and polymyxin, in a single layer or double layer. For these assays seed cultures ATCC 4617 Bordetella bronchiseptica and ATCC 10536 Escherichia coli are recommended.

Technique

The diffusion method for the assay of antibiotics is carried out in accordance with the methodology in the current pharmacopoeias of each country. Scharlau Microbiology's Antibiotic Medium B can be used equally with impregnated paper discs, penicylinders and cut wells as the consistency of the gel is specifically adjusted to suit all of these methodologies.

References

- · ARRET, B.D., P.JOHNSON & A. KIRSCHBAUM (1971) Outline details for Microbiological Assays of Antibiotics: Second revision. J. Pharm. Sci. 60(11):1689-1694
- EUROPEAN PHARMACOPOEIA 7.0 (2011) 7th ed. §. 2.7.2 Microbiological Assay of Antibiotics. EDMH. Council of Europe. Strasbourg.
- ISO/TS 11133-1: 2009. Microbiology of food and animal feeding stuffs.-Guidelines on preparation and production of culture media. Part 1: General guidelines on quality assurance for the preparation of culture media in the laboratory.
- · ISO/TS 11133-2: 2003 Corr. 2004. Microbiology of food and animal feeding stuffs.- Guidelines on preparation and production of culture media. Part 2: Practical guidelines on performance testing of culture media.
- · SANCHO, J., J.GUINEA & R. PARÉS. (1980) Microbiología Analítica Básica. Ed. JIMS. Barcelona.
- · U.S. PHARMACOPOEIA 31 /NATIONAL FORMULARY 26 (2008) Biological Tests and Assays. (81) Antibiotic Microbial Assays. USP Convention Ltd. Rockville, MD.
- · USP 33 NF 28 (2011) <81> Antibiotics Microbial Assays. USP Corp. Inc. Rockville. MD. USA.

Storage

For laboratory use only. Keep tightly closed, away from bright light, in a cool dry place (+4 °C to 30 °C and <60 % RH).

Quality control

Incubation temperature: 35°C ± 2,0

Incubation time: 24 h

Inoculum: 10-100 CFU. Spiral Plate Method (according to standard ISO/TR 11133-1:2000 and ISO/TS 11133-2:2003)

Microorganism	Growth	Remarks
Escherichia coli ATCC 10536	Productivity > 0.70	-
Bordetella bronchiseptica ATCC 4617	Productivity > 0.70	-)