



Reference : 01-444

Scharlau Microbiology - Technical data sheet

Product :
**Yersinia CIN (CEFSULODIN-IRGASAN®-
NOVOBIOCIN) AGAR BASE**

Also known as

CIN Agar; *Yersinia* Selective Agar

Specification

Solid differential medium used for the selective isolation of *Yersinia spp.* from highly polluted samples, according to ISO 10273 standard.

Formula * in g/L

Special peptone.....	20.000	Magnesium sulfate.....	0.010
Yeast extract.....	2.000	Neutral red.....	0.030
Mannitol.....	20.000	Crystal violet.....	0.001
Sodium pyruvate.....	2.000	Agar.....	15.000
Sodium chloride.....	1.000		
Sodium deoxycholate.....	0.500	Final pH 7,4 ±0,2 at 25 °C	

* Adjusted and /or supplemented as required to meet performance criteria

Directions

Suspend 30,25 g in 500 mL of distilled water and bring to the boil. Sterilize in the autoclave at 121°C for 15 minutes. Let it cool to 50-55°C and, aseptically, add the content of a vial of CIN *Yersinia* Selective Supplement (Art. No. 06-143LYO1). Homogenize and pour into plates.

Description

Cefsulodin-Irgasan™-Novobiocin Agar CIN Agar was originally formulated by Schiemann (1979) for detection of *Yersinia enterocolitica*. He subsequently (1982) revised it by substituting sodium deoxycholate for bile salts and reducing the novobiocin content. It relies on the use of selective inhibitory components sodium deoxycholate, crystal violet, cefsulodin, Irgasan® and novobiocin. The basic principle involved is fermentation of mannitol with localised pH reduction which forms a red colony due to the neutral red and a zone of precipitation due to the deoxycholate.

The characteristic appearance of *Yersinia spp.* colonies after an incubation of 18-24 hours at 30°C or 48 hours at 22°C on CIN Agar in air, are round, pink, about 2 mm in diameter with a dark pink centre and surrounded with a precipitation zone. Confirmatory tests are required.

Typical colonies of *Yersinia enterocolitica* will develop as a red bull's-eye surrounded by a transparent border, but will vary considerably among serotypes in colony size, smoothness and the ratio of the border to centre diameter. Most other organisms that are capable of growing on this medium produce larger colonies (> 2 mm in diameter) with diffuse pinkish centres and opaque outer zones. Some strains of *Serratia*, *Citrobacter* and *Enterobacter* on CIN Agar may give a colonial morphology resembling *Yersinia enterocolitica*.

These organisms can be differentiated by simple biochemical tests.

Necessary supplements

Yersinia Selective Supplement (Art. No. 06-143LYO1)

Vial Contents:

Necessary amount for 500 mL of complete medium.

Cefsulodin	7,50 mg
Irgasan®	2,00 mg
Novobiocin	1,25 mg

Distilled water (Solvent)



Reference : 01-444

Scharlau Microbiology - Technical data sheet

Product :

Yersinia CIN (CEFSULODIN-IRGASAN®-NOVOBIOCIN) AGAR BASE

Technique

At present no single isolation procedure is available for the recovery of all pathogenic strains of *Yersinia enterocolitica*. The isolation procedure used will depend on the bio/serogroups of *Yersinia* spp. sought and on the type of sample to be examined. The ISO method for the detection of presumptive pathogenic *Yersinia enterocolitica* includes the parallel use of two isolation procedures:

1. Enrichment in Peptone, Sorbitol and Sels biliaire (PSB) Broth for 2-3 days à 22-25°C with agitation or 5 days without agitation; plating on CIN Agar directly and after alkaline treatment and incubation for 24 hours à 30°C.
2. Enrichment in ITC (Irgasan®-Ticarcillin-Chlorate) Broth for 2 days à 24°C; plating on SSDC (Salmonella-Shigella-Deoxycholate-Calcium Chloride) Agar and incubation for 2 days à 30°C.

Quality control

Incubation temperature: 30 ± 2 °C

Incubation time: 21±3h

Inoculum: Practical range 50- 100 CFU (Productivity) /10⁴-10⁶ CFU (Selectivity) according to ISO 11133:2014/Amd 1:2018

Microorganism	Growth	Remarks
<i>Yersinia enterocolitica</i> ATCC® 9610	Good	-
<i>Escherichia coli</i> ATCC® 25922	Partial inhibition	-
<i>Staphylococcus aureus</i> ATCC® 25923	inhibited	-

References

- ATLAS, R.M. & J.W. SNYDER (1995) Handbook of Media for Clinical Microbiology. CRC Press. Boca Raton. Fla. USA.
- BAYLIS, C.L. (Ed.) (2007) Manual of Microbiological Methods for the Food and Drinks Industry. 5th ed. Guideline No. 43, Campden & Chorleywood Food Research Association. (CCFRA). U.K.
- CORRY, J.E.L., G.D.W. CURTIS & R.M. BAIRD (2003) Handbook of Culture Media for Food Microbiology. Progress in Industrial Microbiology, vol. 37. Elsevier Science Amsterdam.
- De BOER, E. (2003) Isolation of *Yersinia enterocolitica* from foods in "Handbook of Culture Media for Food Microbiology". J.E.L. Corry et al. (Eds.) Elsevier Sci. B.V.
- FDA (Food and Drug Administrations) (1998) Bacteriological Analytical Manual. 8th ed, revision A. AOAC International. Gaithersburg. MD. USA.
- ISENBERG, H.D. (ed.) (1992) Clinical Microbiology Procedures Handbook. ASM. Washington. DC. USA.
- ISO Standard 10273 (2003) Microbiology of food and animal feeding stuffs - Horizontal method for the detection of presumptive pathogenic *Yersinia enterocolitica*.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- SCHIEMAN, D.A. (1979) Synthesis of a selective medium for *Yersinia enterocolitica*. Can. J. Microbiol. 25:1298-1304.
- SCHIEMAN, D.A. (1980) *Yersinia enterocolitica*: Observations on some growth characteristics and response to selective agents. Can. J. Microbiol. 26:1232-1240.
- SCHIEMAN, D.A. (1982) Development of a two step enrichment procedure for recovery of *Yersinia enterocolitica* from food. Appl. Environm. Microbiol. 43:14-27.
- WEAGANT, S.D. & P. FENG (2001) *Yersinia*, in "Compendium of Methods for the Microbiological Examination of Foods". 4th ed. Downes & Ito (Eds.) APHA. Washington. DC. USA.

Storage

For laboratory use only. Keep tightly closed, away from bright light, in a cool dry place (+4 °C to 30 °C).