



Reference: 0BA1019-50

Scharlau Microbiology - Technical Data

Product: Egg Yolk -50ml

Specification

Egg emulsion for microbiological media formulation

Presentation

1 Prepared bottle
Bottles 50 ml
with: 50 ± 3 ml

Packaging Details

1 box with 10 bottle 60 ml. Injectable cap: Plastic screw inner cap. The use of syringes needles with a diameter greater than 0.8 mm is not recommended.

Shelf Life

24 months

Storage

2-14 °C

Composition

Composition (g/l):

Egg Yolk200 ml

Sterile water..... 800 ml

(according to ISO 7932:2004)

Description /Technique

Egg emulsion for different culture media supplementation. Add aseptically to melted bottles of Bacillus cereus base medium cooled to 50°C, before pouring into Petri dishes when cooled to room temperature.

Once solidified on a flat surface, Spread the plates by streaking methodology or by spiral method. Incubate the plates right side up aerobically at 35-37°C for 24-48 hours.

(Incubation times longer than those mentioned above or different incubation temperatures may be required depending on the sample, on the specifications,... This medium can be inoculated directly or after any enrichment broth)

After incubation, enumerate all the colonies that have appeared onto the surface of the agar.

Selective supplementation of the medium suppresses almost all the accompanying flora.

Each laboratory must evaluate the results according to their specifications.

Presumptive isolaton of Bacillus sp must be confirmed by further microbiological and biochemical tests.

Calculate total microbial count per ml of sample by multiplying the average number of colonies per plate by the inverse dilution factor if streaked a diluted sample. Report results as Colony Forming Unit (CFU's) per ml or g along with incubation time and temperature.

Quality control

Physical/Chemical control

Color : yellow

Microbiological control

Add 10 ml of product to 90 ml of Bacillus Cereus Agar base

Inoculate: Practical range 100 ± 20 CFU. min. 50 CFU (productivity)/ 10⁴-10⁶ (selectivity).

Analytical methodology according to ISO 11133:2014/A1:2018; A2:2020.

Aerobiosis. Incubation at 37 °C ± 1, reading after 24-48 ± 2h

Microorganism

Bacillus cereus ATCC® 11778, WDCM 00001

Growth

Good

Sterility control

Inoculate 10 ml of product in 100 ml THIO USP / TSB. Incubate and verify in TSA.

Incubation 48 h at 30-35 °C and 48 h at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.



Reference: 0BA1019-50

Scharlau Microbiology - Technical Data

Product: **Egg Yolk -50ml**

Bibliography

- SMITH, B.A. and BAIRD-PARKER, A.C. (1964) J. Appl. Bact. 27:28.
- VANDERZANT, C. and D.F. SPLITTSTOESSER (1992) Compendium of Methods for the Microbiological Examination of Foods. 3rd. Ed. APHA. Washington.
- leNAN. (1982). Técnicas para el Análisis Microbiológico de Alimentos y Bebidas. Madrid.
- MOSSEL, D.A.A., KOOPMAN, M.J., JONGERIUS, E. (1967) Appl. Microbiol. 15, 650-653.
- NYGREN, B. (1962) Acta Path. Microbiol. Scand. 56, Suppl. 1-160.
- ISO 7932 Standard (2004) 3rd ed. Microbiology of food and animal feeding stuffs. Horizontal method for the enumeration of presumptive *Bacillus cereus*. Colony count technique at 30°C.